

AD-A100 360

TEMPLE UNIV PHILADELPHIA PA SCHOOL OF MEDICINE
GENETICS OF NOVEL HYBRID BACTERIOPHAGE AND DEVELOPMENT OF GENE--ETC(U)
AUG 79 N YAMAMOTO

F/G 6/13

DAMD17-74-C-4051

NL

UNCLASSIFIED

(of)

AD
D-100-360



END
DATE
FILMED
7-81
DTIC

AD A100360

DTIC FILE COPY

AD
II
Annual Progress Report No. 6, (1)

Genetics of Novel Hybrid Bacteriophage and Development of
Generalized Transducing System for Salmonella typhosa

Annual Progress Report
(From 9/1/78 to 3/31/79)

Nobuto/Yamamoto, Ph.D.

August, 1979

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND,

Fort Detrick, Frederick, Maryland 21701

Contract No. DAMD 17-74-C-4051

Temple University
Philadelphia, Pennsylvania 19140

DOD AVAILABILITY STATEMENT: Approved for public release;
distribution unlimited

The findings in this report are not to be construed as an official
Department of the Army position unless so designated by other
authorized documents.

81 6 15 153

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A100360	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Genetics of Novel Hybrid Bacteriophages and Development of Generalized Transducing System for <u>Salmonella typhosa</u>		5. TYPE OF REPORT & PERIOD COVERED Annual Progress Report (From 9/1/78 to 8/31/79)
7. AUTHOR(s) Nobuto Yamamoto		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Temple University School of Medicine 3420 N. Broad Street Philadelphia, Pa. 19140		8. CONTRACT OR GRANT NUMBER(s) DAMD 17-74-C-4051 ✓
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Medical Research and Development Command, Fort Detrick, Frederick, Maryland 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102A 3M161102BS01.00.010
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 23, 1979
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bacteriophage, Hybrid bacteriophage, <u>Salmonella typhimurium</u>, Hybrid bacteria, <u>E. coli-S. typhimurium</u> hybrid, <u>Salmonella typhosa</u>, Generalized transduction, Genetic recombination, Antigen conversion, Cell surface, Gene expression, Bacteriophage tail.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Hybrids phages between Salmonella Phage P22 and coliphage and ϕ80 have been isolated by using <u>E. coli-S. typhimurium</u> hybrids. Among those hybrid phages, ϕ80immP22dis hybrids carrying both immunity related genes, <u>c</u> and <u>Im</u> genes of P22 were isolated. Since P22 tail component gene <u>9</u> and somatic antigen		

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. Abstract (Continued)

conversion gene al are located between the c and Im genes of P22, we examined whether $\phi 80immP22dis$ hybrids carry these genes. Some $\phi 80immP22dis$ hybrids carry gene al but not gene 9 whereas the remaining $\phi 80immP22dis$ hybrids carry gene 9 only. No $\phi 80immP22$ hybrid phages containing both the P22 genes 9 and al were found. These observations suggest that $\phi 80immP22dis$ hybrid is formed as a consequence of multiple crossovers.

Although $\lambda immP22dis$ hybrid phages carry both genes 9 and al, $\phi 80immP22dis$ hybrids carry only one of these genes 9 or al. Since the size of $\phi 80$ phage genome is about 92% of the λ genome, we concluded that the $\phi 80immP22$ gene is unable to contain both genes 9 and al.

Accession For

NTIS	ORNL	X
DTIC	NSA	
ERIC	NSA	
AD	NSA	
AS	NSA	
CS	NSA	
DS	NSA	
ES	NSA	
FS	NSA	
GS	NSA	
HS	NSA	
IS	NSA	
JS	NSA	
KS	NSA	
LS	NSA	
MS	NSA	
NS	NSA	
OS	NSA	
PS	NSA	
QS	NSA	
RS	NSA	
SS	NSA	
TS	NSA	
US	NSA	
VS	NSA	
WS	NSA	
XS	NSA	
YS	NSA	
ZS	NSA	

A

Genetics of Novel Hybrid Bacteriophage and Development of
Generalized Transducing System for Salmonella typhosa

Annual Progress Report
(From 9/1/78 to 8/31/79)

Nobuto Yamamoto, Ph.D.

August, 1979

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND,
FORT DETRICK, FREDERICK, MARYLAND 21701

Temple University
Philadelphia, Pennsylvania 19140

Summary

We mapped the phage chromosomes of hybrids between Salmonella phage P22 and coliphage ϕ 80. Since the genomes of hybrid phages consist of clusters of genes derived from evolutionary diverse bacteriophages, we studied the genetic structure of the homology between parental phages and hybrid phages and formation mechanism of these hybrid phages. In this progress report we showed the origin of genetic segments in the hybrid phage genomes and suggested that the hybrids are formed as a consequence of multiple crossovers.

Foreword

Fundamental studies of viral genetics not only play an important role in increasing our knowledge of the action of viruses in disease processes, but have contributed greatly to our knowledge of the whole problem of cell replication, genetic transfer, gene control, morphogenesis, and antigen conversion. The significance of the study of bacterial hybrids between E. coli and Salmonella has greatly broadened with the recent discoveries of hybrid phage between coliphage and Salmonella phage. The study supported by this contract will bring many important answers for mechanisms of genetic evolution, transduction, recombination, gene expression, antigen conversion and viral replication. In addition, such newly constructed hybrids may prove useful in achieving intergeneric transduction via a hybrid phage vector, of chromosomal genes from different genera of enterobacteriaceae. Therefore such hybrid phages may serve as useful vectors in the genetic engineering of a polyvalent oral attenuated vaccine which expresses immunogenic determinants for antigens of Shigella, Salmonella and perhaps even cholera.

Progress

Present Status of this Project

We have previously reported the isolation of an unusual Salmonella typhimurium hybrid sensitive to coliphage λ and Salmonella phage P22 (Gemski, Baron and Yamamoto, PNAS 69, 3110, 1972). This hybrid, constructed by mating an Escherichia coli-K12 Hfr donor with an S. typhimurium recipient, was characterized as an excellent host for achieving genetic recombination between λ and P22. Two broad hybrid phage classes, each with representative types differing presumably in the extent of gene exchange, have been isolated and described in our previous reports. The λ -P22 hybrid class, which has the protein coat of λ , was found to contain at least the c region of P22. The other class, termed P22- λ , has the protein coat of Phage P22, and has inherited at least the c marker of λ .

By employing an approach similar to that previously used to isolate λ -P22 hybrids, we have been able to isolate hybrids between P22 and coliphage ϕ 80. These newly isolated hybrid phages ϕ 80immP22 were found to be extremely valuable phages for understanding formation mechanism of hybrids between unrelated phages.

1. Isolation and Characterization of Hybrid Phages between E. coli Phage ϕ 80 and Salmonella Phage P22.

E. coli-S. typhimurium hybrid strain WR4027 is a rough bacterium and sensitive to coliphage ϕ 80 for its replication but insensitive to P22 phage because of lack of P22 phage adsorption. Therefore WR4027 lysogenic for phage ϕ 80, WR4027(ϕ 80), is insensitive to P22 phage. By infecting WR4027(ϕ 80) with a mixture of high titer stocks of rough specific Salmonella phages (designated R phages), we were able to isolate R-phage resistant derivatives of WR4027(ϕ 80), designated WR4027(ϕ 80)/R, which are smooth and fully sensitive to P22 phage. Phage P22 stocks grown on this smooth derivative of the ϕ 80 lysogen give rise to recombinants between P22 and ϕ 80. Such recombinants were recovered by plating on a P22 resistant host and immune to ϕ 80, namely WR4027(ϕ 80). They retain the protein coat of ϕ 80 but have acquire the immC region of P22. In addition these ϕ 80immP22 recombinant carries P22 DNA replication genes 12 and 18 as well as the x and erf genes of P22. Some ϕ 80immP22 recombinants, designate ϕ 80immP22dis, contain the immI region as well as the immC region, the two sidely separated loci involved in the bipartite immunity system of P22.

2. Characterization of Unusal Hybrid Phages between E. coli phage ϕ 80 and Salmonella phage P22.

As discussed in the previous report, although λ immP22dis hybrids carry both genes 9 and a1, ϕ 80immP22dis hybrids carry gene 9 or a1 (Fig. 1). As shown in Fig. 2, both λ and ϕ 80 phage genomes contain physically corresponding and functionary similar genes. These pahge genomes also carry genetically inert DNA segments which are located between their respective att and tail (J) genes. However, the entire physical length of ϕ 80 phage genome is about 92% of the size of λ phage genome. This seems to be reflection of difference in sizes of their inert segments: ϕ 80 carries an inert DNA segment smaller than that of λ (Fig. 2). Since the inert segments can be replaced by genes 9 and a1 to form dis hybrid phages,

we concluded that the $\phi 80$ immP22dis hybrid phages are unable to accomodate both genes 9 and a1 simultaneously.

3. Attempts to Isolate Hybrid Phages between P22 and Mutator Phage M_{μ} -1.

Numerous attempts to isolate hybrids between P22 and coli mutator phage M_{μ} -1 were unsuccessful. This may be due to lack of induction of M_{μ} -1 prophage by P22 superinfection although we found that P22 infection of λ or $\phi 80$ lysogens results in induction of their prophages. Dr. Martha Howe supplied us with temperature inducible (ts) mutants of M_{μ} -1 phage. We shall study P22 infection of WR4027 strains lysogenic for these M_{μ} -1 mutants.

Future Research Plan

1. Since we isolated various $\phi 80$ immP22 recombinants, we anticipate finding of P22imm $\phi 80$ recombinants, which carry the early regions, at least the c region, of $\phi 80$ and retain the P22 protein coat.

2. We shall study electron microscopic heteroduplex analyses of $\phi 80$ immP22 hybrid DNA with $\phi 80$ DNA. This study is under progress.

3. We shall also look for recombinants between Salmonella phage P22 and E. coli mutator phage M_{μ} -1.

Publications

Yamamoto, N. Wohlhieter, J.A., Gemski, P. and Baron, L.S. λ immP22dis: A hybrid coliphage λ with both immunity region of Salmonella phage P22, Molecular General Genetics, 166, 233-243, 1978.

Yamamoto, k., Numa, S. Whlhieter, J.A., Gemski, P. and Baron, L.S. Isolation of hybrids between Salmonella phage P22 and coliphage $\phi 80$. Abt. Am. Soc. Microbiol. p. 247, 1979.

Yamamoto, N., Gemski, P. and Baron, L.S. Salmonella somatic O-1 antigen conversion in E. coli K12 carrying Salmonella smooth O-repeating units by λ immP22dis hybrid phages. in preparation.




Figure 1. Genome structures of hybrids between Salmonella phage P22 and coliphage :80.




Figure 2. Genomic structures of P22, λ, :80, λ-P22dis and :80-P22dis phages.

12 Copies

Director (ATTN: SGRD-UWZ-AG)
Walter Reed Army Institute of
Research
Walter Reed Army Medical Center
Washington, D.C. 20012

4 Copies

HQDA (SGRD-AJ)
Fort Detrick
Frederick, Maryland 21701

12 Copies

Defense Documentation Center
ATTN: DDC-TCA
Cameron Station
Alexandria, Virginia 22314

1 Copy

Dean
School of Medicine
Uniformed Services University
of the Health Sciences
4301 Jones Bridge Road
Bethesda, Maryland 20014

1 Copy

Superintendent
Academy of Health Sciences, U.S. Army
ATTN: AHS-COM
Fort Sam Houston, Texas 78234

END

DATE
FILMED

7-8-1

DTIC